

An Examination of LENR Design Improvements, Based on Recently  
Gained Understanding of the LENR Mechanisms:  
A Conventional Explanation of LENR  
Updated 2023

Lynn Bowen, Adjunct Professor, Colorado Mountain College

**Outline**

1. Review of Previous Presentation
2. Neutron scattering and neutron reactions (no neutrons escape)
3. Gamma radiation (very little gamma)
4. Energy to  $^4\text{He}$  ratio
5. Electronic Stopping and Lattice Assisted Nuclear Reactions
6. Experimental ways to increase the probability of a chain reaction

## Review of Previous Talk—Primary and Secondary Reaction in a LENR

❑ Consider a LENR in steady state.

- Two Primary reactions:



- Two Secondary reactions:



- Normal Physics explains it.

- Very little gamma. At the experimentally observed energy.
- No neutrons coming out of the setup.
- The  ${}^4\text{He}$ /energy ratio is correct.

❑ No Weirdness required:

- No unexplained enhancement of  ${}^2\text{d}+{}^2\text{d} \rightarrow {}^4\text{He}+\gamma$  reaction.
- No novel ways needed to overcome the Coulomb barrier.
- No unusual physics is needed at all.

❑ The Mechanism:

- Child products collide with cold  ${}^2\text{d}$
- Frequently and energetically enough.
- Sustains the  ${}^2\text{d}$  to  ${}^2\text{d}$  reaction.
- The conditions in the lattice must be right.

## Review of Previous Talk--Calculations for an Example LENR Reactor

Primary 1 reaction:  ${}^2\text{d}+{}^2\text{d} \rightarrow {}^3\text{t}+\text{p}$

Primary 2 reaction:  ${}^2\text{d}+{}^2\text{d} \rightarrow {}^3\text{He}+\text{n}$

Secondary 1 reaction:  ${}^2\text{d}+{}^3\text{t} \rightarrow {}^4\text{He}+\text{n}$

Secondary 2 reaction:  ${}^2\text{d}+{}^3\text{He} \rightarrow {}^4\text{He}+\text{p}$

- The released energies of the primary and secondary reactions are kinetic energies.
- Eight different child products, total.
- Let's examine a numerical example, using a typical electrolytic set-up in steady state.

### For the Typical Example Reactor:

- 1 watt from the primary reactions
- Reaction rate is  $1.71 \times 10^{12}$  per sec.
  - This is  $8.55 \times 10^{11}$  reactions/second for *each* primary reaction.
  - The palladium core is  $1 \text{ cm}^3$ .
  - The NAE has 10X of 100% loading.
  - Typical electrolytic set-up.
- Charged child products (  ${}^1\text{H}$ ,  ${}^2\text{d}$ ,  ${}^3\text{t}$ ,  ${}^3\text{He}$ , and  ${}^4\text{He}$  ) are trapped in Pd Core.
- Energetic neutrons escape the Pd core.
  - But not the reactor set-up.
  - To be shown in later slide.

## Review of Previous Talk--Calculations for an Example LENR Reactor

- Lengthy and complicated.
  - Conventional physics
- For each child product, calculate:
  - Scattering rate
  - Amount of energized  $^2\text{d}$
  - Average energy.
- The final result is in green on right.
  - $7.4425 \times 10^{12}$  energetic  $^2\text{d}$ 's.
  - Average energy =  $0.5976$  MeV.

| Projectile                            | Source of projectile | initial energy of projectile, in MeV | number of energized D-2 from this projectile | total average energy in MeV for this energized D-2 | number of energized D-2 multiplied by the average energy of the D-2 |
|---------------------------------------|----------------------|--------------------------------------|----------------------------------------------|----------------------------------------------------|---------------------------------------------------------------------|
| Proton                                | Primary1             | 3.0300                               | 3.7008E+11                                   | 0.20895                                            | 7.733E+10                                                           |
| T3                                    | Primary1             | 1.0100                               | 3.1296E+11                                   | 0.19762                                            | 6.185E+10                                                           |
| neutron                               | Primary2             | 2.4525                               | 1.1296E+12                                   | 0.27460                                            | 3.102E+11                                                           |
| He3                                   | Primary2             | 0.8175                               | 1.2099E+12                                   | 0.81396                                            | 9.848E+11                                                           |
| He4                                   | Secondary1           | 3.5180                               | 2.1892E+12                                   | 0.40724                                            | 8.915E+11                                                           |
| neutron                               | Secondary1           | 14.0720                              | 6.1192E+11                                   | 0.29634                                            | 1.813E+11                                                           |
| He4                                   | Secondary2           | 3.6800                               | 1.3611E+12                                   | 0.29295                                            | 3.987E+11                                                           |
| Proton                                | Secondary2           | 14.7200                              | 2.5775E+11                                   | 5.98352                                            | 1.542E+12                                                           |
|                                       |                      |                                      |                                              |                                                    |                                                                     |
|                                       |                      | Sum =                                | 7.4425E+12                                   | Sum =                                              | 4.448E+12                                                           |
|                                       |                      |                                      |                                              |                                                    |                                                                     |
| Average energy of all energized D-2 = |                      |                                      |                                              | 0.5976                                             | MeV                                                                 |

## Is this Enough for a Chain Reaction?

- NAE has very high loading of  $^2\text{d}$ .
- The core volume is  $1 \text{ cm}^3$ .
- Shape is rolled up thin sheet.
- Needed for chain reaction:
  - $1.71 \times 10^{12}$  reactions/sec.
- We have  $1.979 \times 10^{12}$
- Yes! A chain reaction occurs.  
(But, I am ignoring electron stopping.)

| A                                          | B                                                             | C                                                                              | D                                                               | E                                                        | F                                                     | G                                                 | H                                                                              |
|--------------------------------------------|---------------------------------------------------------------|--------------------------------------------------------------------------------|-----------------------------------------------------------------|----------------------------------------------------------|-------------------------------------------------------|---------------------------------------------------|--------------------------------------------------------------------------------|
| Lab energy range of projectile ion, in MeV | Introduction rate of projectile ion in #/cm <sup>3</sup> /sec | Average total distance travelled before the energy drops to next level, in cm. | Reaction cross section at this lab energy, in cm <sup>2</sup> . | Reaction rate at this energy, in #/cm <sup>3</sup> /sec. | The fraction of the lost projectiles due to reaction. | Percentage of the lost projectiles due to escape. | Amount of energetic D-2 remaining in NAE after the energy drops to next level. |
| 0.598 to 0.55                              | 7.4425E+12                                                    | 0.5263                                                                         | 1.61E-25                                                        | 4.275E+11                                                | 0.05744                                               | 1%                                                | 6.9449E+12                                                                     |
| 0.55 to 0.50                               | 6.9449E+12                                                    | 0.5113                                                                         | 1.540E-25                                                       | 3.719E+11                                                | 0.05354                                               | 1%                                                | 6.5073E+12                                                                     |
| 0.50 to 0.45                               | 6.5073E+12                                                    | 0.4671                                                                         | 1.48E-25                                                        | 3.059E+11                                                | 0.04701                                               | 1%                                                | 6.1394E+12                                                                     |
| 0.45 to 0.40                               | 6.1394E+12                                                    | 0.4230                                                                         | 1.39E-25                                                        | 2.455E+11                                                | 0.03998                                               | 1%                                                | 5.8350E+12                                                                     |
| 0.40 to 0.35                               | 5.8350E+12                                                    | 0.3789                                                                         | 1.3E-25                                                         | 1.954E+11                                                | 0.03349                                               | 1%                                                | 5.5831E+12                                                                     |
| 0.35 to 0.30                               | 5.5831E+12                                                    | 0.3349                                                                         | 1.18E-25                                                        | 1.500E+11                                                | 0.02687                                               | 1%                                                | 5.3788E+12                                                                     |
| 0.30 to 0.25                               | 5.3788E+12                                                    | 0.2910                                                                         | 1.06E-25                                                        | 1.128E+11                                                | 0.02098                                               | 1%                                                | 5.2133E+12                                                                     |
| 0.25 to 0.20                               | 5.2133E+12                                                    | 0.2473                                                                         | 8.9E-26                                                         | 7.804E+10                                                | 0.01497                                               | 1%                                                | 5.0839E+12                                                                     |
| 0.20 to 0.15                               | 5.0839E+12                                                    | 0.2041                                                                         | 7.2E-26                                                         | 5.080E+10                                                | 0.00999                                               | 1%                                                | 4.9828E+12                                                                     |
| 0.15 to 0.10                               | 4.9828E+12                                                    | 0.1618                                                                         | 5.2E-26                                                         | 2.851E+10                                                | 0.00572                                               | 1%                                                | 4.9047E+12                                                                     |
| 0.10 to 0.05                               | 4.9047E+12                                                    | 0.1229                                                                         | 3.2E-26                                                         | 1.312E+10                                                | 0.00267                                               | 1%                                                | 4.8427E+12                                                                     |
|                                            |                                                               |                                                                                | Sum =                                                           | 1.979E+12                                                |                                                       |                                                   |                                                                                |

## Neutron Scattering and Neutron Reactions

- Energetic Neutrons **escape** the Pd core.
- **Scatter** in the set-up.
  - High scattering rate
  - In Electrolyte
  - In Thermal water bath.
- If a neutron scatters once, then
  - Scattering cross section is larger.
  - More likely to re-scatter.
  - Quickly **thermalizes**.

The Question is: Can the typical set-up scatter all the energetic neutrons at least once?

Again, let's do the math, using highest energy neutrons (14 MeV) as a worst case.

- One watt of power from primary reactions.
- 97% of the hot t-3 react with cold d-2 (14 MeV neutrons).
- The electrolyte is in a cylindrical beaker, with an inner radius of 2.5 cm, and 25 cm tall.
- There is 0.1 mole, 0.694 gm, of lithium in the heavy water.
- The two borosilicate glass beakers are 1/2 cm and 1 cm, thick respectively.
- The thermal water bath extends 25 cm (about 10") beyond the first beaker.
- The whole set-up has a radius of 28.5 cm x 30 cm tall.

# Neutron Scattering and Neutron Reactions

The scattering rate is calculated for each part, for each element of the set up:

- $^2\text{D}$  and Pd in core.
- $^2\text{D}$ ,  $^{16}\text{O}$ ,  $^6\text{Li}$  in electrolyte
- $^{10}\text{B}$ ,  $^{16}\text{O}$ ,  $^{28}\text{Si}$  in 1st Pyrex
- $^1\text{H}$ ,  $^{16}\text{O}$  in water bath
- $^{10}\text{B}$ ,  $^{16}\text{O}$ ,  $^{28}\text{Si}$  in 2nd Pyrex

The total % probability for one scattering is highlighted in green.

Under these typical conditions, all neutrons are scattered at least once before escaping.

|                                                                        |                                                                                 |                                                                                          |                                                                         |                                                                                                                           |                                                                                |                                                                                |                                                                              |                                                                                                                            |                                                                                                                                      |                                                                                |                                                                                 |                                                                                                                            |                                                                                                                            |                                                                         |
|------------------------------------------------------------------------|---------------------------------------------------------------------------------|------------------------------------------------------------------------------------------|-------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------|--------------------------------------------------------------------------------|------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------|---------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------|
| Inside the Core: the Pd and NAE                                        | Scattering Cross section at this energy with D2 in NAE, in barns. 0.635         | Distance the average neutron travels going through all the NAE's in the core, in cm. 0.1 | Density of D2 in NAE, in atoms/cm <sup>3</sup> 6.800E+23                | Percentage Probability of a neutron scattering with D2 atom at this energy. 4.318                                         | Scattering Cross section at this energy with Pd, in barns. 2.50                | Distance the average neutron travels before leaving the Pd core in cm. 1.3434  | Density of Pd atoms, in atoms/cm <sup>3</sup> 6.803E+22                      | Percentage probability of a neutron scattering with Pd atom at this energy. 22.849                                         | Total percentage of original amount that escapes the core without scattering. 72.833                                                 |                                                                                |                                                                                 |                                                                                                                            |                                                                                                                            | Total % that: escapes the core=72%                                      |
| Inside the heavy water electrolyte: the deuterium, oxygen, and lithium | Scattering Cross section at this energy with D2 in heavy water, in barns. 0.635 | Distance the average neutron travels before leaving the electrolyte, in cm. 2.5          | Density of D2 atoms in electrolyte, in atoms/cm <sup>3</sup> 6.684E+22  | Percentage Probability of original amount, of a neutron scattering with D2 atom in electrolyte, at this energy. 7.729     | Scattering Cross section at this energy with O16 in Electrolyte, in barns. 1.0 | Distance the average neutron travels before leaving the electrolyte in cm. 2.5 | Density of oxygen atoms in electrolyte, in atoms/cm <sup>3</sup> 3.342E+22   | Percentage Probability of original amount, of a neutron scattering with O16 atom in electrolyte, at this energy. 6.086     | Scattering Cross section at this energy with Lithium in Electrolyte, in barns. 0.997                                                 | Distance the average neutron travels before leaving the electrolyte in cm. 2.5 | Density of all Lithium atoms in electrolyte, in atoms/cm <sup>3</sup> 1.230E+20 | Percentage Probability of original amount, of a neutron scattering with Lithium atom in electrolyte, at this energy. 0.022 | Total percentage of original amount that escaped both the Pd core and the electrolyte without scattering or reacting. 62.5 | escapes both core and electrolyte = 63%                                 |
| Inside the first borosilicate glass beaker                             | Scattering Cross section at this energy with boron in Pyrex, in barns. 0.815    | Distance the average neutron travels before leaving the Pyrex, in cm. 0.5                | Density of all boron atoms in Pyrex, in atoms/cm <sup>3</sup> 5.588E+21 | Percentage Probability of original amount, of a neutron scattering with Boron atom in Pyrex, at this energy. 0.143        | Scattering Cross section at this energy with O16 in Pyrex, in barns. 0.5       | Distance the average neutron travels before leaving the Pyrex in cm. 1         | Density of all oxygen atoms in Pyrex, in molecules/cm <sup>3</sup> 1.223E+22 | Percentage Probability of original amount, of a neutron scattering with O16 atom in electrolyte, at this energy. 0.384     | Scattering Cross section at this energy with Si28 in Pyrex, in barns. 0.720                                                          | Distance the average neutron travels before leaving the Pyrex in cm. 0.5       | Density of all Silicon atoms in Pyrex, in atoms/cm <sup>3</sup> 8.780E+21       | Percent Probability of original amount, of a neutron scattering with Si atom in electrolyte, at this energy. 0.198         | Total percentage of original amount that escapes the core, electrolyte, and pyrex without scattering or reacting. 62.0     | escapes core, electrolyte, and 1st pyrex = 62%                          |
| Inside the thermal water bath                                          | Scattering Cross section at this energy with H1 in water bath, in barns. 0.688  | Distance the average neutron travels before leaving the water bath, in cm. 25            | Density of H1 atoms in water bath, in atoms/cm <sup>3</sup> 6.691E+22   | Percentage Probability of original amount, of a neutron scattering with H1 atom in the water bath, at this energy. 71.699 | Scattering Cross section at this energy with O16 in water bath, in barns. 1.0  | Distance the average neutron travels before leaving the water bath, in cm. 25  | Density of O16 atoms in water bath, in atoms/cm <sup>3</sup> 2.974E+22       | Percentage Probability of original amount, of a neutron scattering with O16 atom in the water bath, at this energy. 46.317 | Total percentage of original amount that escapes the core, electrolyte, pyrex, and waterbath without scattering or reacting. -11.224 |                                                                                |                                                                                 |                                                                                                                            |                                                                                                                            | escapes core, electrolyte, pyrex, and water bath = -11%                 |
| Inside the second borosilicate glass beaker                            | Scattering Cross section at this energy with boron in Pyrex, in barns. 0.815    | Distance the average neutron travels before leaving the Pyrex, in cm. 1                  | Density of all boron atoms in Pyrex, in atoms/cm <sup>3</sup> 5.588E+21 | Percentage Probability of original amount, of a neutron scattering with Boron atom in Pyrex, at this energy. -0.051       | Scattering Cross section at this energy with O16 in Pyrex, in barns. 1.0       | Distance the average neutron travels before leaving the Pyrex in cm. 0.5       | Density of all oxygen atoms in Pyrex, in molecules/cm <sup>3</sup> 1.223E+22 | Percentage Probability of original amount, of a neutron scattering with O16 atom in electrolyte, at this energy. -0.069    | Scattering Cross section at this energy with Si28 in Pyrex, in barns. 0.720                                                          | Distance the average neutron travels before leaving the Pyrex in cm. 1         | Density of all Silicon atoms in Pyrex, in atoms/cm <sup>3</sup> 8.780E+21       | Percentage Probability of original amount, of a neutron scattering with Si atom in electrolyte, at this energy. -0.0710    | Total percentage of original amount that escapes the core, electrolyte, and pyrex without scattering or reacting. -11.25   | escapes core, electrolyte, 1st pyrex, water bath and 2nd pyrex = -11.2% |

## Neutron Scattering and Neutron Reactions

❑ For Thermal Neutrons, these four nuclear reactions will absorb them. Notice, more  $^4\text{He}$  and heat are created.



- The reaction rate is not enough to change isotopic ratio.



- Cross section of **941** barns.
- Used in hot fusion reactors
  - Stop neutrons
  - Generate heat.



Cross section is **3840** barns.

Used as neutron moderator in fission reactors.

Contained in any borosilicate glass (Pyrex or Dewars) in the set-up.



- High density of  $^1\text{H}$  in water bath.
- Used to shield neutrons in fission reactors.

❑ In a typical set-up, how many of the thermal neutrons will react, and how many escape?

## Neutron Scattering and Neutron Reactions

The thermal neutron reaction rate is calculated for every part, in every element of the set up:

- $^2\text{D}$  and  $\text{Pd}$  in the core.
- $^2\text{D}$ ,  $^{16}\text{O}$ ,  $^6\text{Li}$  in the electrolyte
- $^{10}\text{B}$ ,  $^{16}\text{O}$ ,  $^{28}\text{Si}$  in the 1st Pyrex beaker
- $^1\text{H}$ ,  $^{16}\text{O}$  in the water bath
- $^{10}\text{B}$ ,  $^{16}\text{O}$ ,  $^{28}\text{Si}$  in the 2nd Pyrex beaker

The total % probability of a thermal neutron reaction is shown in green.

Under these typical conditions, all thermal neutrons react and do not escape.

|                                                                         |                                                                                               |                                                                     |                                                                 |                                                                |                                                                        |                                                                                                        |                                                                        |                               |                                                                        |                                                                      |                                                                               |                               |                                                                         |                                                  |
|-------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|---------------------------------------------------------------------|-----------------------------------------------------------------|----------------------------------------------------------------|------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------|-------------------------------|------------------------------------------------------------------------|----------------------------------------------------------------------|-------------------------------------------------------------------------------|-------------------------------|-------------------------------------------------------------------------|--------------------------------------------------|
| In Pd core and NAE                                                      |                                                                                               |                                                                     |                                                                 |                                                                |                                                                        |                                                                                                        |                                                                        |                               |                                                                        |                                                                      |                                                                               |                               |                                                                         | Total % that reacts in the core = 0.00155        |
| Volume of the NAE for the total sample, in cm <sup>3</sup>              | Distance the average neutron travels before leaving the NAE's, going through the core, in cm. | Density of D2 in NAE, in number per cm <sup>3</sup>                 | Neutron to D2 reaction cross section at this energy, in barns.  | Neutron to D2 reaction rate, per second per cm <sup>3</sup> .  | Fractional probability of a neutron reacting with a D2 at this energy  | Volume of the Pd lattice in the core for the total sample, not including the NAE, in cm <sup>3</sup> . | Distance the average neutron travels before leaving the Pd core in cm. | Density of Pd-Natural in Core | Neutron to Pd-Natural reaction cross section at this energy, in barns. | Neutron to Pd-Natural reaction rate                                  | Fractional probability of a neutron reacting with a Pd-Natural at this energy |                               |                                                                         |                                                  |
| 0.60                                                                    | 0.2                                                                                           | 6.80E+23                                                            | 5.00E-04                                                        | 8.71E+02                                                       | 5.15E-10                                                               | 2.00                                                                                                   | 1.26E+00                                                               | 6.80E+22                      | 7.17E+00                                                               | 2.62E+07                                                             | 1.55E-05                                                                      |                               |                                                                         |                                                  |
| In electrolyte                                                          |                                                                                               |                                                                     |                                                                 |                                                                |                                                                        |                                                                                                        |                                                                        |                               |                                                                        |                                                                      |                                                                               |                               |                                                                         | Total % that reacts in the electrolyte = 0.02799 |
| Volume of the electrolyte for the total sample, in cm <sup>3</sup>      | Distance the average neutron travels before leaving the electrolyte, in cm.                   | Density of D2 in electrolyte, in number per cm <sup>3</sup>         | Neutron to D2 reaction cross section at this energy, in barns.  | Neutron to D2 reaction rate, per second per cm <sup>3</sup> .  | Fractional probability of a neutron reacting with a D2 at this energy  | Density of O16 in electrolyte, in number per cm <sup>3</sup>                                           | Neutron to O16 reaction cross section at this energy, in barns.        | Neutron to O16 reaction rate  | Fractional probability of a neutron reacting with a O16 at this energy | Density of Li6 in electrolyte, in number per cm <sup>3</sup>         | Neutron to Li6 reaction cross section at this energy, in barns.               | Neutron to Li6 reaction rate  | Fractional probability of a neutron reacting with a Li6 at this energy  |                                                  |
| 491                                                                     | 5                                                                                             | 6.68E+22                                                            | 5.00E-04                                                        | 1.75E+06                                                       | 1.03453E-06                                                            | 3.34E+22                                                                                               | 2.00E-04                                                               | 3.50E+05                      | 2.06906E-07                                                            | 9.57E+18                                                             | 9.40E+02                                                                      | 4.72E+08                      | 2.79E-0                                                                 |                                                  |
| In borosilicate glass surrounding electrolyte                           |                                                                                               |                                                                     |                                                                 |                                                                |                                                                        |                                                                                                        |                                                                        |                               |                                                                        |                                                                      |                                                                               |                               |                                                                         | Total % that reacts in the 1st Pyrex = 1.16704   |
| Volume of the borosilicate glass around electrolyte, in cm <sup>3</sup> | Distance the average neutron travels before leaving the borosilicate glass, in cm.            | Density of B10 in borosilicate glass, in number per cm <sup>3</sup> | Neutron to B10 reaction cross section at this energy, in barns. | Neutron to B10 reaction rate, per second per cm <sup>3</sup> . | Fractional probability of a neutron reacting with a B10 at this energy | Density of O16 in borosilicate glass, in number per cm <sup>3</sup>                                    | Neutron to O16 reaction cross section at this energy, in barns.        | Neutron to O16 reaction rate  | Fractional probability of a neutron reacting with a O16 at this energy | Density of Si28 in borosilicate glass, in number per cm <sup>3</sup> | Neutron to Si28 reaction cross section at this energy, in barns.              | Neutron to Si28 reaction rate | Fractional probability of a neutron reacting with a Si28 at this energy |                                                  |
| 215.98                                                                  | 1                                                                                             | 1.11E+21                                                            | 3.85E+03                                                        | 1.98E+10                                                       | 1.17E-02                                                               | 1.22E+22                                                                                               | 2.00E-04                                                               | 1.13E+04                      | 6.65E-09                                                               | 8.78E+21                                                             | 1.80E-01                                                                      | 7.29E+06                      | 4.31E-0                                                                 |                                                  |
| Light water in water bath                                               |                                                                                               |                                                                     |                                                                 |                                                                |                                                                        |                                                                                                        |                                                                        |                               |                                                                        |                                                                      |                                                                               |                               |                                                                         | Total % that reacts in the water bath = 82.79477 |
| Volume of the light water in water bath, in cm <sup>3</sup>             | Distance the average neutron travels before leaving the borosilicate glass, in cm.            | Density of H1 in light water, in number per cm <sup>3</sup>         | Neutron to H1 reaction cross section at this energy, in barns.  | Neutron to H1 reaction rate, per second per cm <sup>3</sup> .  | Fractional probability of a neutron reacting with a B10 at this energy | Density of O16 in light water in water bath, in number per cm <sup>3</sup>                             | Neutron to O16 reaction cross section at this energy, in barns.        | Neutron to O16 reaction rate  | Fractional probability of a neutron reacting with a O16 at this energy |                                                                      |                                                                               |                               |                                                                         |                                                  |
| 73183.40                                                                | 40                                                                                            | 6.69E+22                                                            | 3.35E-01                                                        | 1.40E+12                                                       | 8.28E-01                                                               | 2.97E+22                                                                                               | 2.00E-04                                                               | 3.71E+08                      | 2.19E-04                                                               |                                                                      |                                                                               |                               |                                                                         |                                                  |
| Borosilicate glass around light water                                   |                                                                                               |                                                                     |                                                                 |                                                                |                                                                        |                                                                                                        |                                                                        |                               |                                                                        |                                                                      |                                                                               |                               |                                                                         | Total % that reacts in the 2nd Pyrex = 58.05589  |
| Volume of the borosilicate glass around light water, in cm <sup>3</sup> | Distance the average neutron travels before leaving the borosilicate glass, in cm.            | Density of B10 in borosilicate glass, in number per cm <sup>3</sup> | Neutron to B10 reaction cross section at this energy, in barns. | Neutron to B10 reaction rate, per second per cm <sup>3</sup> . | Fractional probability of a neutron reacting with a B10 at this energy | Density of O16 in borosilicate glass, in number per cm <sup>3</sup>                                    | Neutron to O16 reaction cross section at this energy, in barns.        | Neutron to O16 reaction rate  | Fractional probability of a neutron reacting with a O16 at this energy | Density of Si28 in borosilicate glass, in number per cm <sup>3</sup> | Neutron to Si28 reaction cross section at this energy, in barns.              | Neutron to Si28 reaction rate | Fractional probability of a neutron reacting with a Si28 at this energy |                                                  |
| 5372.12                                                                 | 2                                                                                             | 1.11E+21                                                            | 3.85E+03                                                        | 9.83E+11                                                       | 5.80E-01                                                               | 1.22E+22                                                                                               | 2.00E-04                                                               | 5.60E+05                      | 3.31E-07                                                               | 8.78E+21                                                             | 1.80E-01                                                                      | 3.63E+08                      | 2.14E-0                                                                 |                                                  |
| Sum of reaction probabilites, in % = 142.05                             |                                                                                               |                                                                     |                                                                 |                                                                |                                                                        |                                                                                                        |                                                                        |                               |                                                                        |                                                                      |                                                                               |                               |                                                                         |                                                  |

## Gamma Radiation from the Set-up

Gamma is due to thermal neutron absorption:  ${}^A(X) + n \rightarrow {}^{A+1}(X) + \gamma$

To calculate the gamma radiation:

- Determine the number of thermal neutrons. **(Already Done.)**
- Calculate the number of thermal neutron absorption reactions per second. **(Already Done.)**
- Determine energy and intensity of gamma. **(Done below.)**
- And how much gamma is absorbed by the set-up. **(Done below.)**

| Reaction | Energy of gamma radiation in MeV | Initial Intensity of the gamma radiation in gamma rays per second | Post intensity of the gamma radiation after penetrating 25 cm of water | Estimated post-intensity of the gamma radiation after penetrating 1 cm of borosilicate glass | Post intensity of this gamma after passing through both water and borosilicate glass. |
|----------|----------------------------------|-------------------------------------------------------------------|------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| D2+n     | 6.26                             | 1.23E+06                                                          | 1.99E+05                                                               | 1.13E+06                                                                                     | 1.82E+05                                                                              |
| Pd105+n  | 9.56                             | 1.85E+07                                                          | 2.98E+06                                                               | 1.69E+07                                                                                     | 2.72E+06                                                                              |
| O16+n    | 4.14                             | 2.62E+08                                                          | 4.23E+07                                                               | 2.40E+08                                                                                     | 3.87E+07                                                                              |
| Si28+n   | 8.473                            | 2.46E+08                                                          | 3.97E+07                                                               | 2.25E+08                                                                                     | 3.63E+07                                                                              |
| H1+n     | 2.22                             | 9.87E+11                                                          | 2.46E+11                                                               | 9.02E+11                                                                                     | 2.25E+11                                                                              |
|          |                                  |                                                                   |                                                                        | Sum=                                                                                         | 2.25E+11                                                                              |

➤ Note the numbers in green. These dominate.

➤ This model predicts a small amount of gamma at 2.22 MeV.

➤ This is seen experimentally.

## Excess Heat Compared to $^4\text{He}$

|                             | <u>Heat Energy Released</u> | <u>Amount of <math>^4\text{He}</math> Produced</u>                    |
|-----------------------------|-----------------------------|-----------------------------------------------------------------------|
| <u>Primary reactions:</u>   | 1 watt.                     | No $^4\text{He}$ is produced.                                         |
| <u>Secondary reactions:</u> | 3.39 watts                  | $8.421 \times 10^{11} + 3.480 \times 10^{11} = 11.901 \times 10^{11}$ |
| <u>Totals:</u>              | 4.39 watts                  | $1.1901 \times 10^{12}$                                               |

Ratio:  $(4.39 \text{ joules/sec}) / (1.1901 \times 10^{12} \text{ } ^4\text{He/sec}) = 3.7 \times 10^{-12} \text{ joules per } ^4\text{He}$

**= 23.05 MeV per  $^4\text{He}$ .**

- This energy is right on target.
  - Not coincidence.
  - Going from two  $^2\text{D}$  to  $^4\text{He}$ , a NET release of energy, no matter what path is taken.
  - Simple conservation of energy.
- There is **more  $^4\text{He}$**  and **more energy** created in the lithium and boron.
  - Experimentally, the exact ratio has slight **variations**—it **depends on the set-up**.

## Electron Stopping Power

Thus, this model answers the questions, explains the experimental data, gives the right values.  
No miracles. No unexplainable behaviors. No violations of thermodynamics.

😞 **Except for one thing... Electron Stopping Power.**

😊 This came to my attention last year.

😊 I thought this would reduce the reaction by a factor of 8 or so.

😊 I thought changing the shape of the core could accommodate this.

- Changing the shape works, but not enough.

😊 I thought I could reduce the excess heat output.

- Excess heat is not a factor for the chain reaction.

💀 Bottom Line: Electron Stopping power kills the reaction.

The formula for electron stopping  $S$  as a function of velocity  $v$  is:

$$S(v) = \frac{4\pi n}{m_e v^2} Z^2 k_e^2 e^4 \log \left( \frac{2m_e v^2}{I} \right)$$

where  $I$  is the average excitation energy--experimentally determined for each material.

## Electron Stopping Power and LANR

What are some possible ways to reduce or mitigate Electron Stopping Power?

- Apply a counter electric field to motivate the ions forward.
- Use a superconducting material as the core.
- Reduce the electron coefficient of heat.
- Apply a strong magnetic field.
- Increase the temperature.
- Increase the current.
- Ionization.

Is there something is occurring **in the lattice**, maybe more than one thing, that allows the subsequent  $^2\text{d}$ -to- $^2\text{d}$  reactions to occur?

After all, we DO call the phenomenon:

**Lattice Assisted Nuclear Reactions.**

## Electron Stopping and LANR

### Interesting facts about the PdD<sub>x</sub> lattice:

#### ➤ Superconductivity

- 😊 PdD<sub>x</sub> is a superconductor.
  - 😬 An unconventional high-temperature superconductor.
  - 😬 It's behavior can't be explained by any theory. (Sound familiar?)
  - 😬 The experimental data for its superconducting behavior is not always reproducible. (Sound familiar?)
- 😊 Theory for conventional superconductors tells us that electrons pair up
  - 😬 To form "Cooper pairs"
  - 😬 Due to an unknown undefined unexplained attractive force. (Seriously.)
- 😊 Superconductivity is strongly related to current density and magnetism.
  - 😬 Superconductivity must have a critical current density,  $J_c$ .
  - 😬 When superconducting, the normal magnetic field caused by a current flowing in a wire, goes away.

## Electron Stopping and LANR

### Interesting facts about the PdD<sub>x</sub> lattice:

- PdD<sub>x</sub> has *ionized* <sup>2</sup>d in it.
  - The <sup>2</sup>d is metallic hydrogen.
  - This accounts for the high absorption of <sup>2</sup>d in Pd.
- The electrons in PdD<sub>x</sub> do not follow rules for filling electron shells.
  - The electrons in Pd fill a higher level d-shell before filling the lower s-shell.
  - The β-phase of PdD<sub>x</sub> is characterized by s-electrons filling the d-band.
  - Complexities of the electron quantum mechanics.
- Magnetic Susceptibility
  - The magnetic susceptibility of PdD<sub>x</sub> varies significantly with x.
  - PdD<sub>x</sub> changes from being paramagnetic to diamagnetic as x increases
  - Going from alpha to beta phase.

## Electron Stopping and LANR

### More Interesting Facts about the PdD<sub>x</sub> lattice:


➤ There are 2 Heat Coefficients: Atomic and Electronic.

➤ The Electronic Heat Coefficient is  $\gamma$ , gamma.

- $\gamma$  is dependent on the electron density of states.
- For pure Pd,  $\gamma = 9.5 \text{ mJ}/(\text{mol} \cdot \text{K}^2)$ .
- For PdD<sub>x</sub>,  $\gamma$  is 6x smaller than for pure Pd.
- The electrons vibrate more.
- This is the superconducting region for PdD<sub>x</sub>.

➤ The <sup>2</sup>D to <sup>2</sup>D reaction curve.

- It is not a cliff-like drop-off at low energies.
- Rather, a gradual sloped drop-off.
- Due to tunneling (or other effect) overcoming
- the Coulomb barrier.
- (The Electromagnetic Model explains this. See my poster.)



## **Electron Stopping and LANR**

### **Interesting facts about Superconductors**

When excited ions are traveling inside a lattice, there are two ways that defects are created:

- Elastic displacement.
  - Collision by charged ionic particles with other nuclei.
  - Related to nuclear stopping,  $S_n$ .
- Electron excitation.
  - Collision by charged ionic particles with the electrons.
  - Related to electron stopping,  $S_e$ .
- For ion projectiles with an energy below 2 MeV
  - Experimental evidence implies
  - Superconductors are dominated by nuclear stopping
  - Not electron stopping.

Quoting from the literature...

## Electron Stopping and LANR

### Interesting facts about Superconductors

Ishikawa et al. found that for low energy projectiles ( $\leq 2\text{MeV}$ ), the defect production rate in a superconductor is proportional to the nuclear stopping power, rather than electron stopping.

of the  $c$ -axis lattice parameter with increasing fluence for all irradiations. The slope of  $c$ -axis lattice parameter against fluence, which corresponds to the defect production rate, is separated into two contributions; the effect via elastic displacement and the effect via electronic excitation. The former contribution exhibits a linear increase against the nuclear stopping power,  $S_n$ . The latter contribution is scaled by the primary ionization rate,  $dJ/dx$ , rather than by the electronic stopping power,  $S_e$ , and is nearly proportional to  $(dJ/dx)^4$ .

the value of  $(\Delta c/c_0)/\Phi$ . For low energy ( $\leq 2\text{ MeV}$ ) ion irradiation,  $(\Delta c/c_0)/\Phi$ , which corresponds to the defect production rate, is proportional to the nuclear stopping power,  $S_n$ . On the other hand, for high energy (80 MeV–3.80 GeV) ion irradiation, the contribution of electronic excitation to

## Electron Stopping and LANR

### Interesting facts about Superconductors

From Barbour et al.:

“For ions with energies less than several MeV, the dominant mechanism causing irradiation-induced degradation of  $T_c$  is collisional damage.”

In other words, the damage in the lattice is created by collisions, which implies  $S_n$  is the dominant stopping power.

This is in agreement with previous experiments:

“The defect production rate, which is obtained from the initial slope of the resistance–fluence curve, is almost proportional to the nuclear stopping power,  $S_n$ . The  $T_c$  depression due to irradiation is also nearly proportional to  $S_n$ . These results indicate that the defect production process is dominated by elastic displacement. .... In high- $T_c$  superconductor  $\text{EuBa}_2\text{Cu}_3\text{O}_y$  irradiated with 2 MeV electrons and 0.5–2.0 MeV ions, the normal-state resistance change and  $T_c$  depression due to irradiation are nearly proportional to  $S_n$ , indicating that the elastic displacement is a dominant damage production process.”

In other words, these other behaviors of the lattice that follow this same trend,  $S_n$  dominates.

**There is something going on in the lattice that mitigates the electron stopping power.**

## **Experimental ways to Increase the Probability of a Chain Reaction:**

- **What can we change experimentally to build a better reactor?**
  - The  $^2d$  density doesn't actually play a strong role for the chain reaction.
    - Has to be high enough for the NAE to be formed.
    - Has to be high enough to put the PdD<sub>x</sub> lattice is in the super conduction region.
  - Size and shape of the Pd core matter--but to a limited extent.
- **Other things may not directly affect the reaction rate, but they may affect the lattice, the superconductivity, and the electron stopping. CANR.**
  - Current density. Applied voltages. Oscillations of these.
  - Temperature.
  - Magnetism.
  - Homogeneity of the loading.
  - Pressure.

## Experimental ways to Increase the Probability of a Chain Reaction:

### **Use a starter**

- Especially for well-shielded set-up
- Could cause intermittency and inconsistency
- Alpha emitter, abutting the core.

### **Lithium Li-6 and Boron B-10 are good.**

- Reduce emission of stray neutrons.
- Pyrex. Dewar. Electrolyte.
- $^4\text{He}$  particles come out of electrolyte
- Neutrons were formed and reacted, to create the  $^4\text{He}$ .

### **The shape needs to be right.**

- One that:
  - Moderates
  - Utilizes
  - Contains the neutrons.
- Excellent shapes, with large surface area.
  - Bundle of thin wires
  - Rolled-up thin foil
  - Porous collection of small Pd grains.

If a neutron escapes the Pd core, it is moderated, scattered, and re-enters another part of the Pd core.

The more we contain the neutrons inside the core and use their energy, the better.

## Summary and Conclusions

**All the so-called “miracles” of LENR are explained.**

- ✓ The Coulomb barrier is overcome by the transference of kinetic energy to the cold  $^2\text{d}$ .
- ✓ Neutrons are emitted from the reactions, but they do not escape the set-up.
- ✓ Very little gamma radiation occurs, because only the tertiary reactions create them.
- ✓ Gamma occurs at 2.22 MeV, as is verified experimentally.
- ✓ The predicted energy per  $^4\text{He}$  ratio is on target. Not coincidental.

No unusual or novel physics is needed. Simply apply conventional nuclear physics to the LENR.

### **EXCEPT...**

- **...Except** for electron stopping.
- We already know there are unexplained behaviors of superconductivity and the electrons.
- We already know there is something unusual with the electrons occurring in the lattice.
- We don't know exactly WHAT it is.

Further research is needed, both theoretically and experimentally, to clarify exactly what is happening.

## Conclusions

- A conventional explanation of LENR should not be casually rejected.
- That is an unfortunate assumption, made before actually studying the facts.
- The conventional explanation can explain more experimental behavior than other models.
- These concepts need to be more thoroughly investigated.

Thank you for your attention.

